

Table 10.1

## Kaposi Sarcoma

## Number of Cases

By SEER Location and Year of Diagnosis, 1975-2011

## All Races, Males, All Ages

## Year of Diagnosis

| SEER Location                           | 1975-1979 | 1980-1981 | 1982-1983 | 1984-1985 | 1986-1987 | 1988-1989 | 1990-1991 | 1992-1993 | 1994-1995 | 1996-1997 | 1998-1999 | 2000-2001 | 2002-2003 | 2004-2005 | 2006-2007 | 2008-2009 | 2010-2011 |
|---|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|
| SEER 9 <sup>b</sup>                     | 134       | 79        | 325       | 888       | 1,573     | 2,131     | 2,231     | 2,057     | 1,800     | 841       | 392       | 402       | 336       | 372       | 334       | 343       | 342       |
| SEER 9 except SF-Oakland                | 102       | 44        | 85        | 261       | 428       | 776       | 841       | 976       | 874       | 464       | 245       | 265       | 223       | 245       | 237       | 250       | 247       |
| San Francisco-Oakland SMSA <sup>c</sup> | 32        | 35        | 240       | 627       | 1,145     | 1,355     | 1,390     | 1,081     | 926       | 377       | 147       | 137       | 113       | 127       | 97        | 93        | 95        |
| San Francisco City/County               | 15        | 21        | 200       | 539       | 953       | 1,076     | 1,089     | 792       | 689       | 254       | 85        | 82        | 72        | 83        | 62        | 61        | 65        |

Age-Adjusted Cancer Incidence Rates<sup>a</sup>

By SEER Location and Year of Diagnosis, 1975-2011

## All Races, Males, All Ages

## Year of Diagnosis

| SEER Location                           | 1975-1979 | 1980-1981 | 1982-1983 | 1984-1985 | 1986-1987 | 1988-1989 | 1990-1991 | 1992-1993 | 1994-1995 | 1996-1997 | 1998-1999 | 2000-2001 | 2002-2003 | 2004-2005 | 2006-2007 | 2008-2009 | 2010-2011 |
|---|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|
| SEER 9 <sup>b</sup>                     | 0.4       | 0.5       | 1.7       | 4.2       | 7.1       | 9.2       | 9.3       | 8.2       | 7.0       | 3.2       | 1.5       | 1.5       | 1.3       | 1.4       | 1.2       | 1.3       | 1.2       |
| SEER 9 except SF-Oakland                | 0.4       | 0.4       | 0.6       | 1.6       | 2.3       | 4.0       | 4.2       | 4.6       | 4.0       | 2.1       | 1.1       | 1.2       | 1.0       | 1.1       | 1.1       | 1.1       | 1.1       |
| San Francisco-Oakland SMSA <sup>c</sup> | 0.7       | 1.3       | 7.0       | 17.2      | 30.0      | 34.0      | 34.0      | 25.6      | 22.0      | 9.0       | 3.5       | 3.1       | 2.7       | 3.0       | 2.3       | 2.1       | 2.1       |
| San Francisco City/County               | 1.1       | 3.1       | 25.0      | 65.2      | 111.2     | 124.9     | 125.9     | 89.1      | 77.4      | 28.3      | 9.4       | 8.9       | 8.0       | 9.2       | 6.8       | 6.6       | 6.9       |

<sup>a</sup> Rates are per 100,000 males and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).<sup>b</sup> The SEER 9 areas are San Francisco-Oakland SMSA, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah and Atlanta.<sup>c</sup> The San Francisco-Oakland SMSA represents 15% of all persons in the SEER areas. For this age group, all SEER areas may not be generalizable to the United States for Kaposi Sarcoma.

- Statistic not shown. Rate based on less than 4 cases for the time interval.